UTAH - WEEKLY INFLUENZA TEXT SUMMARY

Influenza Activity in Utah:

Laboratory confirmed influenza has been detected in Utah. This includes 514 hospitalizations, the majority in high risk categories. 71% of hospitalizations have been identified as Type A, 21% Type B, and 8% unknown. Influenza-like illness levels are decreasing.

As of 5/17/2008, 1 pediatric influenza-associated deaths has been reported in Utah for the current season.

National influenza surveillance data can be found at www.cdc.gov/flu. Please note, states determine for themselves how to estimate weekly influenza activity. The classification system used in Utah primarily serves to provide a geographical description of how influenza is circulating in the state and not season severity. Utah is currently reporting "sporadic" activity.

This website will be updated every Wednesday at 1 PM.

INFLUENZA PREVENTION

How to protect yourself and others:

1. Utahns should get an influenza vaccine. **Contact your provider or the Immunization Hotline (1-800-275-0659) to locate an influenza shot. 2. All Utahns should practice "respiratory etiquette" to reduce the spread of respiratory diseases.

- Stay away from other people when you are sick.
- ** Cover your mouth and nose with a disposable tissue when you cough or sneeze. Throw the tissue away
- ** Wash your hands.
- 3. We urge schools and employers to be supportive of these preventive measures.

Influenza Activity for Current Season and Past Three Seasons

Figure 1. Percentage of visits for influenza-like illness (ILI) reported by sentinel providers- Utah, 2007-2008 and previous three influenza seasons







